Necessary and sufficient condition on the fixed-energy scattering data for the potential to be spherically symmetric

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1989 Inverse Problems 5 L45

(<http://iopscience.iop.org/0266-5611/5/5/001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 194.95.184.111

The article was downloaded on 29/06/2011 at 08:24

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Necessary and sufficient condition on the fixed-energy scattering data for the potential to be spherically symmetric

A G Ramm† and O L Weaver‡

† Mathematics Department, Kansas State University, Manhattan, KS 66506, USA

‡ Strahlenzentrum, University of Giessen, 6300 Giessen, Federal Republic of Germany, and Physics Department, Kansas State University, Manhattan, KS 66506, USA

Received 20 June 1989

Abstract. Consider the Schrödinger equation for fixed energy and a real-valued compactly supported potential $q(x) \in L^2_{\text{loc}}$, $x \in \mathbb{R}^3$. Let $A(\theta', \theta)$ be the corresponding scattering amplitude for incident and exit directions θ and θ' respectively. It is proved that the potential is spherically symmetric if and only if $A(\theta', \theta)$ for this single energy depends only on the angle between θ and θ' , that is $A(\theta', \theta) = A(\theta' \cdot \theta)$ for all $\theta', \theta \in S^2$, the unit sphere. Thus, if this condition holds at a single energy $\kappa^2 > 0$ it holds for all $\kappa > 0$ under the above assumption about $q(x)$.

Let

$$[\nabla^2 + \kappa^2 - q(x)]u = 0 \quad \text{in } \mathbb{R}^3 \quad \kappa > 0 \text{ is fixed.} \quad (1)$$

Without loss of generality let us assume in what follows that $\kappa = 1$. The scattering solution is defined as the solution to (1) with the asymptotics

$$u = \exp(i\theta \cdot x) + A(\theta', \theta)r^{-1} \exp(ir) + o(r^{-1}) \quad \text{as } r = |x| \rightarrow \infty, \quad x r^{-1} = \theta'. \quad (2)$$

The function $A(\theta', \theta)$ is the scattering amplitude. Let us assume

$$q \in Q_a := \{q: q(x) = \bar{q}(x) \in L^2(B_a), q(x) = 0 \text{ in } B'_a\} \quad (3)$$

where $B_a = \{x: |x| \leq a\}$, $B'_a := \mathbb{R}^3 \setminus B_a$. If $q(x) = q(r)$, $r = |x|$, then it is well known that

$$A(\theta', \theta) = A(\theta' \cdot \theta) \quad \forall \theta', \theta \in S^2 \quad (4)$$

where S^2 is the unit sphere in $\mathbb{R}^3$. The basic result of this letter, theorem 1, has been formulated as a conjecture in [1].

Theorem 1. Assume that (3) holds. Then $q(x) = q(r)$ iff (4) holds. Thus (4) holds at all energies iff it holds at one.

The result of this type in the case when $A(\theta', \theta, \kappa) = A(\theta' \cdot \theta, \kappa)$ for all $\theta', \theta \in S^2$ and all $\kappa > 0$ has been proved in [1]. In [2–4] a theory for inverse scattering problems with fixed-energy data is given and in [5–6] one finds basic properties of the scattering amplitude. Our proof is very short and is based on the uniqueness theorem proved in [2–4, 7, 8].

Proof. First we formulate the uniqueness theorem from [4].

Theorem 2. If $q_j \in Q_a$, $j = 1, 2$, and $A_1(\theta', \theta) = A_2(\theta', \theta)$ for all $\theta', \theta \in S^2$ then $q_1 = q_2$. Here $A_j(\theta', \theta)$ is the scattering amplitude corresponding to q_j , $j = 1, 2$, at a fixed $\kappa > 0$.

Assume that (4) holds. Our basic idea is to show that (4) implies

$$A_q(\theta', \theta) = A_{q \circ R}(\theta', \theta) \quad \forall \theta', \theta \in S^2 \text{ and } \forall R \in O(3) \quad (5)$$

where $O(3)$ is the group of rotations in $\mathbb{R}^3$. Here $A_q(\theta', \theta)$ is the scattering amplitude corresponding to the potential $q(x)$, and $q \circ R$ is the potential whose value at a point x is $(q \circ R)(x) := q(R^{-1}x)$. Since $q(R^{-1}x) \in Q_a$ if $q(x) \in Q_a$, it follows from theorem 2 and equation (5) that

$$q(x) = q(R^{-1}x) \quad \forall R \in O(3). \quad (6)$$

This means that $q(x) = q(|x|) = q(r)$. Thus the proof is complete as long as (5) is established. Equation (5) follows from the equations

$$A_{q \circ R}(R\theta', R\theta) = A_q(\theta', \theta) \quad (7)$$

and

$$A_q(\theta', \theta) = A_q(R\theta', R\theta). \quad (8)$$

Equation (7) holds for any $q(x)$ and is an immediate consequence of the definition of the scattering amplitude $A_q(\theta', \theta)$. Equation (8) follows from our basic assumption (4), since $\theta' \cdot \theta = R\theta' \cdot R\theta$ for any $R \in O(3)$.

Let us explain (7) in detail. Let $A_q(\theta', \theta)$ be the scattering amplitude in a certain coordinate system, say τ . Consider the same scattering amplitude in a coordinate system τ' in which any vector which in the coordinate system τ was x is equal to Rx , where $R \in O(3)$. In the coordinate system τ' the potential q evaluated at a point y has the value $(q \circ R)(y) = q(R^{-1}y)$. For example, suppose that the potential q evaluated at a point x in the coordinate system τ has the value $q(x) = d \cdot x$, where d is a constant vector and the dot denotes inner product. Then, in the coordinate system τ' , vector d is equal to Rd , so that the potential q evaluated at a vector y in the coordinate system τ' is equal to $Rd \cdot y = d \cdot R^{-1}y = q(R^{-1}y) := (q \circ R)(y)$. Here we use the fact that $R' = R^{-1}$, where R' is the transposed transformation; that is, the fact that rotations in $\mathbb{R}^3$ are orthogonal transformations. Thus, in the coordinate system τ' the value of the scattering amplitude $A_q(\theta', \theta)$ is equal to $A_{q \circ R}(R\theta', R\theta)$. This is the desired equation (7). Theorem 1 is proved.

Remark 1. Our argument shows that, for any class of potentials $q(x)$ for which the uniqueness theorem for the inverse scattering problem holds for certain data, condition (4) is necessary and sufficient for $q(x)$ to be spherically symmetric if this condition holds for the above data. For example, if (4) holds for all sufficiently large $\kappa > 0$ then $q(x) = q(|x|)$ if q belongs to the class of potentials for which the uniqueness of the solution of the inverse scattering problem is proved for the data $A(\theta', \theta, \kappa)$ given for all $\theta', \theta \in S^2$ and all κ sufficiently large. These are potentials $q(x)$ for which $|q(x)| \leq c(1 + |x|)^{-b}$, $b > 3$, $c = \text{constant} > 0$; see [6] (for an even larger class of potentials with $b > 1$, see [9]).

The argument in this paper is applicable to the scattering by obstacles such as, for example, in the problem considered in [1, 7, 10].

AGR thanks NSF, ONR and SFB 256 and OLW thanks the Fulbright Commission for support.

References

- [1] Ramm A G 1989 Necessary and sufficient condition for a scattering amplitude to correspond to a spherically symmetric scatterer *Appl. Math. Lett.* **2**
- [2] Ramm A G 1988 Multidimensional inverse problems and completeness of the products of solutions to PDE *J. Math. Anal. Appl.* **134** 211–53
- [3] Ramm A G 1988 Uniqueness theorems for multidimensional inverse problems with unbounded coefficients *J. Math. Anal. Appl.* **136** 568–74
- [4] Ramm A G 1988 Recovery of potential from the fixed-energy scattering data *Inverse Problems* **4** 877–86
- [5] Ramm A G 1986 *Scattering by Obstacles* (Dordrecht: Reidel)
- [6] Ramm A G 1987 Characterization of the scattering data in multidimensional inverse scattering problem *Inverse problems: An Interdisciplinary Study* ed P C Sabatier (New York: Academic) pp 153–67
- [7] Ramm A G 1989 Completeness of the products of solutions to PDE and inverse problems *Int. Conf. on Inverse Problems (Bulgaria, September 1989)*, Preprint SFB 256, Bonn
- [8] Ramm A G 1989 Multidimensional inverse scattering problems and completeness of the products of solutions to homogeneous PDE *Z. Angew. Math. Mech.* **69** T13–22
- [9] Saito Y 1986 Asymptotic and approximate formulas in the inverse scattering problem for the Schrödinger operator *Lecture Notes in Mathematics* **1218** (Berlin: Springer) pp 190–200
- [10] Ramm A G 1989 Symmetry properties for scattering amplitudes and applications to inverse problems *Preprint SFB 256*, Bonn